

Optimizing Enzymatic Maceration of Blackcurrant Mash by Response Surface Methodology

Krasimira Petrova¹, Todorka Petrova¹, Petya Ivanova¹, Milena Ruskova¹, Ivan Bakalov¹, Kiril Mihalev², and Nikolay Penov²

1. Food Research and Development Institute, Bulgaria

2. University of Food Technologies, Bulgaria

Abstract: Pre-press maceration with commercial enzyme preparation (Pectinex Ultra Color) was investigated in experimental blackcurrant (Omota variety) juice production using response surface methodology (RSM). Enzyme dosage and maceration time were varied at a constant temperature of 50°C and the total polyphenols (TPP), total monomeric anthocyanins (TMA), antioxidant capacity and juice yields (Y) were compared. The contents of total polyphenols and anthocyanins ranged from 828.57 to 1274.96 mg GAE/100 g and 326.76 to 383.49 mg CGE/100 g blackcurrant juice, respectively. The total antioxidant capacity determined by the free radical scavenging activity (DPPH) of the juice ranged from 3977.82 to 4580.18 $\mu\text{mol TE}/100\text{ g}$ and ferric reducing antioxidant power (FRAP) from 4749.19 to 6579.74 $\mu\text{mol TE}/100\text{ g}$. Juice yields ranged from 44.57 to 50.31% by wet weight of the mash. The criterion established to determine the optimal conditions of enzymatic maceration of blackcurrant mash was to find the conditions leading to high values of total monomeric anthocyanins, total polyphenols, and juice yield.

Key words: response surface methodology (RSM), enzymatic maceration, juice, blackcurrant

1. Introduction

Blackcurrants (*Ribesnigrum* L.) are one of the most popular and widely cultivated species in Europe. It is traditionally used for producing of juices, nectars, jams, and nutraceutical ingredients. Blackcurrants are rich in phenolic compounds, notably anthocyanins and hydroxycinnamic acids. The water-soluble anthocyanins in black currants are mainly found in the skin [1].

Pectolytic enzymes are commonly used in industrial blackcurrant juice production to facilitate juice extraction [2]. These enzymes are used during conventional processing cause degradation of the cell wall matrix and increased the juice yield [3]. Enzyme-assisted blackcurrant juice processing significantly increases the contents of anthocyanins

and other phenolic compounds in the juice [4].

An optimal central composite design (OCCD) is an experimental approach to define empirical models or equations for describing the effect of test variables and their interactions on the respective responses [5].

Response surface methodology (RSM) is an effective tool which uses quantitative data in an experimental design to optimize a process [6]. RSM has been used for optimizing processes in fruit and vegetable juice production [5, 7, 8].

The aim of this study was to optimize the enzymatic mash maceration using RSM, thus allowing extraction of polyphenol-rich blackcurrant juice.

2. Material and Methods

2.1 Chemicals

For the analytical purposes DPPH [2,2-diphenyl-1-picrylhydrazyl]; TPTZ [2,4,6-Tris(2-pyridyl)-s-triazine]; Trolox [(±)-6-Hydroxy-2,5,7,8-tetramethylchroman

Corresponding author: Krasimira Petrova, Ph.D., research area: food technologies. E-mail: kراسи_petrova80@abv.bg.

-2-carboxylic acid] (Sigma-Aldrich, Steinheim, Germany); Folin-Ciocalteu's (FC) reagent (Merck, Darmstadt, Germany); and Gallic acid monohydrate (Fluka, Buchs, Switzerland) were used. All the other reagents and solvents used were of analytical grade.

2.2 Enzyme Preparation

Pectinex Ultra Color was obtained from Novozymes A/S (Bagsvaerd, Denmark). It is produced from *Aspergillusaculeatus* and *Aspergillusniger*, with declared activity of 10 000 PECTU/mL, and is brown liquid. The suitable maceration temperature is 50°C and is active in the pH range of 3.0-5.0.

2.3 Plant Material and Processing

Blackcurrant (*Ribesnigrum* L.) fruits, at a commercial maturity stage (harvest 2015) were from Research institute of mountain stockbreeding and agriculture, Troyan, Bulgaria.

After sorting to eliminate unripened, overripened or damaged fruits, blackcurrant fruits were frozen stored until used.

Frozen fruits were processed according to the flow diagram shown in Fig. 1.




Fig. 1 Process flowchart for extraction of the blackcurrant juice.

2.4 Analytical Methods

All measurements were performed with a Helios Omega UV-Vis spectrophotometer equipped with VISIONlite software (all from Thermo Fisher Scientific, Madison, WI, USA) using 1 cm path length cuvettes.

The contents of total polyphenols (TPP) and total monomeric anthocyanins (TMA) were determined by the method of Singleton and Rossi [9] and the pH-differential method [10], respectively, modified as described by Dinkova et al. [11].

The total antioxidant capacity was determined by the DPPH (free radical scavenging activity) and FRAP (ferric reducing antioxidant power) assay, following the methods of Brand-Williams et al. [12] and Benzie and Strain [13], respectively, with some modifications [11].

2.5 Experimental Design and Data Analysis

The results reported in the present study are the mean values of at least three determinations, and the coefficients of variation, expressed as the percentage ratio between the standard deviations and the mean values, were found to be < 5% in all cases.

An optimal central composite design (OCCD) was used to investigate the effect of the enzyme dosage (X_1) and maceration time (X_2) on the response (y) in 11 runs [6]. The outline of the experimental design is outlined in Table 1.

Table 1 Experimental design for the enzymatic maceration.

Run №	Coded levels		Actual levels	
	X_1	X_2	Enzyme dosage (%E/S ^a)	Maceration time (min)
1	-1	-1	0.02	30
2	+1	-1	0.18	30
3	-1	+1	0.02	210
4	+1	+1	0.18	210
5	-1	0	0.02	120
6	+1	0	0.18	120
7	0	-1	0.10	30
8	0	+1	0.10	210
9	0	0	0.10	120
10	0	0	0.10	120
11	0	0	0.10	120

^a ml enzyme preparation per 100 g substrate.

A second order polynomial model for the dependent variables TMA, TPP, DPPH, FRAP, and juice yield (response, y) was established to fit the experimental data:

$$y = b_0 + \sum_{i=1}^n b_i x_i + \sum_{i=1}^n b_{ii} x_i^2 + \sum_{i=1}^n \sum_{j=1}^n b_{ij} x_i x_j \quad (1)$$

where b_0 = intercepts, b_i are linear, b_{ii} are quadratic, and b_{ij} are interaction regression coefficient terms.

SYSTAT statistical software (SPSS Inc., Chicago, USA, version 7.1) and Excel were used to analyze the data results.

3. Results and Discussion

The mean values of TMA, TPP, DPPH, FRAP, and juice yield are shown in Table 2. The contents of total polyphenols and anthocyanins ranged from 828.57 to 1274.96 mg GAE/100 g and 326.76 to 383.49 mg CGE/100 g blackcurrant juice, respectively. The total antioxidant capacity determined by the free radical scavenging activity (DPPH) of the juice ranged from 3977.82 to 4580.18 $\mu\text{mol TE}/100\text{ g}$ and ferric reducing antioxidant power (FRAP) from 4749.19 to 6579.74 $\mu\text{mol TE}/100\text{ g}$. Juice yields ranged from 44.57 to 50.31% by wet weight of the mash.

Table 2 TMA, TPP, DPPH, FRAP, and blackcurrant juice yield.

Run №	TMA ^a	TPP ^b	DPPH ^c	FRAP ^c	Yield (%)
1	326.76	828.57	3977.82	4749.19	44.57
2	366.78	893.32	3944.34	5178.07	47.58
3	383.49	1138.27	4330.31	6288.62	46.00
4	370.90	1274.96	4580.18	6579.74	46.19
5	353.34	966.50	3979.66	5988.87	44.87
6	369.94	1128.12	4231.05	6182.20	45.20
7	355.76	1017.01	3971.74	5181.64	50.31
8	379.33	1259.28	4715.08	6547.19	50.09
9	375.87	1136.29	4270.88	6244.37	49.60
10	375.32	1172.40	4298.94	6434.10	49.41
11	367.64	1086.43	4358.94	6102.71	49.12

^a Results are presented as mg cyanidin 3-glucoside equivalents (CGE) per 100 g juice;

^b Results are presented as mg gallic acid equivalents (GAE) per 100 g juice;

^c Results are presented as μmol trolox equivalents (TE) per 100 g juice.

The regression equations describing the effect of enzymatic maceration of blackcurrant mash on the TMA, TPP, DPPH, FRAP, and juice yield are given in Table 3. The resulting equations with high accuracy describe the change of the contents of the dependent variable ($R^2 > 0.9$). The coefficients in the regression equations can be used to examine the significance of each term relative to each other when used with coded values [14].

Each of the estimated effects and interactions are shown in the standardized diagram — the Pareto diagram (Fig. 2). It consists of horizontal blocks with lengths proportional to the absolute values of the estimated effects, divided by their standard errors. The vertical line in the Pareto diagram represents the value of the Student criterion at 95% confidence level and separates factors that are significant to those that are not. The diagrams show the predominance of the maceration time (factor B) for TMA, TPP, DPPH, and FRAP. The quadratic effect due to the enzyme dosage (factor A) has the highest impact on the juice yield.

The effect of changes in enzyme dosage and maceration time on TMA, TPP, DPPH, FRAP, and juice yield is given in Fig. 3. The values of anthocyanins, total polyphenols and antioxidant capacity increased with an increase in the maceration time. The yield increased between 0.02 and 0.12% E/S and then decreased between 0.12 and 0.18% E/S. Similar results were reported by Dinkova et al. [15].

Table 3 Coefficients of regression models for TMA, TPP, DPPH, FRAP, and blackcurrant juice yield.

Term	TMA	TPP	DPPH	FRAP	Yield
	Coefficients				
constant	304.72	727.25	3793.15	4092.97	42.54
X_1	569.75*	3422.45*	5041.55	7787.34*	151.05*
X_2	0.41*	1.60*	0.14*	20.04*	-0.02
X_1X_1	-1294.04*	-14829.1*	-26238.2*	-26552.8	-659.75*
X_1X_2	-1.83*	2.50	9.84	-4.78	-0.10*
X_2X_2	-0.0003	-0.0005	0.009	-0.05*	0.0001*
R^2	0.97	0.96	0.95	0.98	0.98

X_1 – enzyme dosage (%E/S), X_2 – maceration time (min), R^2 – coefficient of determination.

*Significant at 95% CI.




Fig. 2 Estimated effects of regression model coefficients on TMA (a), TPP (b), DPPH (c), FRAP (d), and juice yield (e).




Fig. 3 Effect of enzyme dosage and maceration time on TMA (a), TPP (b), DPPH (c), FRAP (d), and juice yield (e)

The criterion established to determine the optimal conditions of enzymatic maceration of blackcurrant mash was to find the conditions leading to high values of total monomeric anthocyanins (TMA > 370 mg CGE/100 g), total polyphenols (TPP > 1100 mg GAE/100 g), and juice yield (Y > 48%). Optimization was carried out by the superposition of the contour plots for predicted TMA, TPP, and juice yield. The optimum area for the enzymatic mash maceration is presented on Fig. 4 (the darkened area). On each point of this area corresponds conditions of enzymatic maceration of blackcurrant mash leading to obtain the juice with high values of TMA, TPP, and Y.

4. Conclusions

The effect of conditions of enzymatic maceration of blackcurrant mash on the total monomeric anthocyanins, total polyphenols, antioxidant capacity, and juice yield was investigated. The choice of the preferred conditions was according to survey of other researchers on enzymatic maceration. The linear effect due to the maceration time had the highest impact on TMA, TPP, DPPH, and FRAP. The quadratic effect due to the enzyme dosage had the highest impact on the juice yield. The criterion established to determine




Fig. 4 Graphical optimization of the enzymatic maceration of blackcurrant mash.

the optimal conditions of enzymatic maceration of blackcurrant mash was to find the conditions leading to high values of total monomeric anthocyanins, total polyphenols, and juice yield. Maceration time from 120 to 210 min and enzyme dosage from 0.06 to 0.16% E/S could be recommended as the optimal conditions of enzymatic maceration of blackcurrant mash for obtaining juice with high values of TMA, TPP, and Y.

References

- [1] A. Landbo and A. Meyer, Effects of different enzymatic maceration treatments on enhancement of anthocyanins and other phenolics in black currant juice, *Innovative Food Science and Emerging Technologies* 5 (2004), 503-513.
- [2] O. Laaksonen, L. Mäkilä, R. Tahvonen, H. Kallio and B. Yang, Sensory quality and compositional characteristics of blackcurrant juices produced by different processes, *Food Chemistry* 138 (2013) 2421-2429.
- [3] H. Hilz, E. Bakx, H. Schols and A. Voragen, Cell wall polysaccharides in black currants and bilberries — Characterization in berries, juice, and press cake, *Carbohydrate Polymers* 59 (2005) 477-488.
- [4] R. Bagger-Jørgensen and A. Meyer, Effects of different enzymatic pre-press maceration treatments on the release of phenols into blackcurrant juice, *European Food Research and Technology* 219 (2004) 620-629.
- [5] Y. Sun, Z. Wang, J. Wu, F. Chen, X. Liao and X. Hu, Optimizing enzymatic maceration in pretreatment of carrot juice concentrate by response surface methodology, *International Journal of Food Science and Technology* 41 (2006) 1082-1089.
- [6] A. Lambrev, *Bases of the Engineering Experiment in the Investigation of Machines and Devices for the Food Industry*, “Ch. G. Danov”, Plovdiv, 1994, p. 211. (in Bulgarian)
- [7] P. Rai, G. Majumdar, S. Dasgupta and S. De, Optimizing pectinase usage in pretreatment of mosambijuce for clarification by response surface methodology, *Journal of Food Engineering* 64 (2004) 397-403.
- [8] H. Sin, S. Yusof, Sheikh, N. Hamid and R. Rahman, Optimization of enzymatic clarification of sapodilla juice using response surface methodology, *Journal of Food Engineering* 73 (2006) 313-319.
- [9] V. Singleton and J. Rossi, Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents, *American Journal of Enology and Viticulture* 50 (1963) 3828-3834.
- [10] M. Giusti and R. Wrolstad, Anthocyanins: Characterization and measurement with UV visible spectroscopy, in: R. E. Wrolstad (Ed.), *Current Protocols*

- in *Food Analytical Chemistry*, New York: Wiley, Unit F1.2.1-13, 2001.
- [11] R. Dinkova, V. Shikov, K. Mihalev, Z. Velchev, H. Dinkov and P. Mollov, Changes in the total anthocyanins and polyphenols during processing of wild berries into freshly pressed juices, *Journal of EcoAgri Tourism* 8 (2012) (1) 254-259.
- [12] W. Brand-Williams, M. Cuvelier and C. Berst, Use of a free radical method to evaluate antioxidant activity, *Lebensmittel Wissenschaft and Technologie* 28 (1995) 25-30.
- [13] I. Benzie and J. Strain, The Feric reducing ability of plasma (FRAP) as a measure of antioxidant activity: The FRAP assay, *Analytical Biochemistry* 189 (1996) 70-76.
- [14] N. Rastogi and K. Rashmi, Optimization of enzymatic liquefaction of mango pulp by response surface methodology, *European Food Research and Technology* 209 (1999) 57-62.
- [15] R. Dinkova, V. Shikov, K. Mihalev, Z. Velchev, N. Penov, H. Dinkov and P. Mollov, Optimizing enzymatic maceration of bilberry mash by response surface methodology, *Journal of Mountain Agriculture on the Balkans* 16 (2013) (1) 327-339.